

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 88113448.0

(51) Int. Cl.⁴: **C07D 215/14 , C07D 215/18 ,
C07D 215/20 , C07D 405/06 ,
A61K 31/47**

(22) Date of filing: 18.08.88

(30) Priority: 20.08.87 JP 207224/87
26.01.88 JP 15585/88
03.08.88 JP 193606/88

(43) Date of publication of application:
22.02.89 Bulletin 89/08

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: **NISSAN CHEMICAL INDUSTRIES LTD.**
7-1, 3-chome Kanda-Nishiki-cho
Chiyoda-ku Tokyo(JP)


(72) Inventor: Fujikawa, Yoshihiro Nissan
Chemical Industries Ltd. Chuo Kenkyusho,
722-1
Tsuboi-cho Funabashi-shi Chiba-ken(JP)
Inventor: Suzuki, Mikio Nissan
Chemical Industries Ltd. Chuo Kenkyusho,
722-1
Tsuboi-cho Funabashi-shi Chiba-ken(JP)
Inventor: Iwasaki, Hiroshi Nissan
Chemical Industries Ltd. Chuo Kenkyusho,
722-1
Tsuboi-cho Funabashi-shi Chiba-ken(JP)
Inventor: Sakashita, Mitsuaki Nissan
Chemical Industries Ltd
Seibutsukagaku Kenkyusho, 1470
Oaza-shiraoka
Shiraoka-machi Minamisaitama-g
Saitama-k(JP)
Inventor: Kitahara, Masaki Nissan Chemical
Industries Ltd.
Seibutsukagaku Kenkyusho, 1470
Oaza-shiraoka
Shiraoka-machi Minamisaitama-g
Saitama-k(JP)

(74) Representative: Wächtershäuser, Günter, Dr.
Tal 29
D-8000 München 2(DE)


EP 0 304 063 A2

(54) Quinoline type mevalonolactones.


(57) A compound of the formula:



wherein R^1 , R^2 , R^3 , R^4 and R^5 are independently hydrogen, C_{1-6} alkyl, C_{1-6} cycloalkyl, C_{1-3} alkoxy, n -butoxy, i -butoxy, sec -butoxy, R^7R^8N - (wherein R^7 and R^8 are independently hydrogen or C_{1-3} alkyl), trifluoromethyl, trifluoromethoxy, difluoromethoxy, fluoro, chloro, bromo, phenyl, phenoxy, benzyloxy, hydroxy, trimethylsilyloxy, diphenyl- t -butylsilyloxy, hydroxymethyl or $-O(CH_2)_tOR^{19}$ (wherein R^{19} is hydrogen or C_{1-3} alkyl, and t is 1, 2 or 3); or when located at the ortho position to each other, R^1 and R^2 , or R^3 and R^4 together form $-CH=CH-CH=CH-$; or when located at the ortho position to each other, R^1 and R^2 together form $-OC(R^{15})(R^{16})O-$ (wherein R^{15} and R^{16} are independently hydrogen or C_{1-3} alkyl); Y is $-CH_2-$, $-CH_2CH_2-$, $-CH=CH-$, $-CH_2-CH=CH-$ or $-CH=CH-CH_2-$; and Z is $-Q-CH_2WCH_2CO_2R^{12}$,



(wherein Q is $-C(O)-$, $-C(OR^{13})_2-$ or $-CH(OH)-$; W is $-C(O)-$, $-C(OR^{13})_2-$ or $-C(R^{11})(OH)-$; R^{11} is hydrogen atom or C_{1-3} alkyl; R^{12} is hydrogen or R^{14} (wherein R^{14} is physiologically hydrolyzable alkyl or M (wherein M is NH_4 , sodium, potassium, $1/2$ calcium or a hydrate of lower alkyl amine, di-lower alkyl amine or tri-lower alkyl amine)); two R^{13} are independently primary or secondary C_{1-6} alkyl; or two R^{13} together form $-(CH_2)_2-$ or $-(CH_2)_3-$; R^{17} and R^{18} are independently hydrogen or C_{1-3} alkyl; and R^{19} is hydrogen, C_{1-6} alkyl, C_{2-3} alkenyl, C_{3-6} cycloalkyl,



(wherein R^9 is a hydrogen atom, C_{1-4} alkyl, C_{1-3} alkoxy, fluoro, chloro, bromo or trifluoromethyl), phenyl- $(CH_2)_m-$ (wherein m is 1, 2 or 3), $-(CH_2)_nCH(CH_3)-phenyl$ or phenyl- $(CH_2)_nCH(CH_3)-$ (wherein n is 0, 1 or 2).

QUINOLINE TYPE MEVALONOLACTONES

The present invention relates to novel mevalonolactones having a quinoline ring, processes for their production, pharmaceutical compositions containing them and their pharmaceutical uses particularly as anti-hyperlipidemic, hypolipoproteinemic and anti-atherosclerotic agents, and intermediates useful for their production and processes for the production of such intermediates.

5 Some fermentation metabolic products such as compactine, CS-514, Mevinolin or semi-synthetic derivatives or fully synthetic derivatives thereof are known to be inhibitors against HMG-CoA reductase which is a rate limiting enzyme for cholesterol biosynthesis. (A. Endo J. Med Chem., 28(4) 401 (1985))


CS-514 and Mevinolin have been clinically proved to be potentially useful anti-hyperlipoproteinemic agents, and they are considered to be effective for curing or preventing diseases of coronary artery sclerosis or atherosclerosis. (IXth Int. Symp. Drugs Affect. Lipid Metab., 1986, p30, p31, p66)

However, with respect to fully synthetic derivatives, particularly hetero aromatic derivatives of inhibitors against HMG-CoA reductase, limited information is disclosed in the following literatures:


WPI ACC NO. 84-158675, 86-028274, 86-098816, 86-332070, 87-124519, 87-220987, 88-07781, 88-008460, 88-091798 and 88-112505.

15 The present inventors have found that mevalonolactone derivatives having a quinoline ring, the corresponding dihydroxy carboxylic acids and salts and esters thereof have high inhibitory activities against cholesterol biosynthesis wherein HMG-CoA reductase acts as a rate limiting enzyme. The present invention has been accomplished on the basis of this discovery.


The novel mevalonolactone derivatives of the present invention are represented by the following formula 20 I:



wherein R^1 , R^2 , R^3 , R^4 and R^6 are independently hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{1-3} alkoxy, *n*-butoxy, *i*-butoxy, *sec*-butoxy, R^7R^8N - (wherein R^7 and R^8 are independently hydrogen or C_{1-3} alkyl), trifluoromethyl, trifluoromethoxy, difluoromethoxy, fluoro, chloro, bromo, phenyl, phenoxy, benzyloxy, hydroxy, trimethylsilyloxy, diphenyl-*t*-butylsilyloxy, hydroxymethyl or $-O(CH_2)_lOR^{19}$ (wherein R^{19} is hydrogen or C_{1-3} alkyl, and l is 1, 2 or 3); or when located at the ortho position to each other, R^1 and R^2 , or R^3 and R^4 together form $-CH=CH-CH=CH-$; or when located at the ortho position to each other, R^1 and R^2 together form $-OC(R^{15})(R^{16})O-$ (wherein R^{15} and R^{16} are independently hydrogen or C_{1-3} alkyl); Y is $-CH_2-$, $-CH_2CH_2-$, $-CH=CH-$, $-CH_2-CH=CH-$ or $-CH=CH-CH_2-$; and Z is $-Q-CH_2WCH_2-CO_2R^{12}$, 40



(wherein Q is -C(O)-, -C(OR¹³)₂- or -CH(OH)-; W is -C(O)-, -C(OR¹³)₂- or -C(R¹¹)(OH)-; R¹¹ is hydrogen or C₁₋₃ alkyl; R¹² is hydrogen or R¹⁴ (wherein R¹⁴ is physiologically hydrolyzable alkyl or M (wherein M is NH₄, sodium, potassium, 1/2 calcium or a hydrate of lower alkylamine, di-lower alkylamine or tri-lower alkylamine)); two R¹³ are independently primary or secondary C₁₋₆ alkyl; or two R¹³ together form -(CH₂)₂- or -(CH₂)₃-; R¹⁷ and R¹⁸ are independently hydrogen or C₁₋₃ alkyl; and R⁵ is hydrogen, C₁₋₆ alkyl, C₂₋₃ alkenyl, C₃₋₆ cycloalkyl,



(wherein R⁹ is hydrogen, C₁₋₄ alkyl, C₁₋₃ alkoxy, fluoro, chloro, bromo or trifluoromethyl), phenyl-(CH₂)_m- (wherein m is 1, 2 or 3), -(CH₂)_nCH(CH₃)-phenyl or phenyl-(CH₂)_nCH(CH₃)- (wherein n is 0, 1 or 2).

Various substituents in the formula I will be described in detail with reference to specific examples. However, it should be understood that the present invention is by no means restricted by such specific examples.

C₁₋₆ alkyl for R¹, R², R³, R⁴, R⁶ and R⁸ includes, for example, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl and t-butyl. C₁₋₃ alkoxy for R¹, R², R³, R⁴ and R⁶ includes, for example, methoxy, ethoxy, n-propoxy and i-propoxy.

C₁₋₃ alkyl for R¹¹ includes, for example, methyl, ethyl, n-propyl and i-propyl.

C₁₋₃ alkyl for R¹³ includes, for example, methyl, ethyl, n-propyl and i-propyl.

Alkyl for R¹⁴ includes, for example, methyl, ethyl, n-propyl, i-propyl, n-butyl and i-butyl.

M is a metal capable of forming a pharmaceutically acceptable salt, and it includes, for example, sodium and potassium.

CO₂M includes, for example, -CO₂NH₄ and -CO₂H⁺ (primary to tertiary lower alkylamine such as trimethylamine).

C₁₋₆ alkyl for R⁵ includes, for example, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, n-pentyl and n-hexyl.

C₃₋₆ cycloalkyl for R⁵ includes, for example, cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

C₂₋₃ alkenyl for R⁵ includes, for example, vinyl and i-propenyl.

Phenyl-(CH₂)_m- for R⁵ includes, for example, benzyl, β-phenylethyl and γ-phenylpropyl.

Phenyl-(CH₂)_nCH(CH₃)- for R⁵ includes, for example, α-phenylethyl and α-benzylethyl.

C₁₋₃ alkyl for R⁷ and R⁸ includes, for example, methyl, ethyl, n-propyl and i-propyl.

Further, these compounds may have at least one or two asymmetric carbon atoms and may have at least two to four optical isomers. The compounds of the formula I include all of these optical isomers and all of the mixtures thereof.

Among compounds having carboxylic acid moieties falling outside the definition of -CO₂R¹² of the carboxylic acid moiety of substituent Z of the compounds of the present invention, those which undergo physiological hydrolysis, after intake, to produce the corresponding carboxylic acids (compounds wherein the -CO₂R¹² moiety is -CO₂H) are equivalent to the compounds of the present invention.

Now, preferred substituents of the compounds of the present invention will be described.

In the following preferred, more preferred still further preferred and most preferred examples, the numerals for the positions of the substituents indicate the positions on the quinoline ring. For example, N' shown by e.g. 1' or 2' indicates the position of the substituent on the phenyl substituted at the 4-position of the quinoline ring (the carbon connected to the quinoline ring is designated as 1'). The meanings of the respective substituents are the same as the above-mentioned meanings.

Preferred substituents for R¹, R² and R⁶ are hydrogen, fluoro, chloro, bromo, C₁₋₃ alkyl, C₁₋₃ alkoxy, C₃₋₆ cycloalkyl, dimethylamino, hydroxy, hydroxymethyl, hydroxyethyl, trifluoromethyl, trifluoromethoxy, difluoromethoxy, phenoxy and benzyloxy.

Further, when R⁶ is hydrogen, it is preferred that R¹ and R² together form methylenedioxy.

As preferred examples for R³ and R⁴, when R⁴ is hydrogen, R³ is hydrogen, 3'-fluoro, 3'-chloro, 3'-methyl, 4'-methyl, 4'-chloro and 4'-fluoro.

Other preferred combinations of R³ and R⁴ include 3'-methyl-4'-chloro, 3',5'-dichloro, 3',5'-difluoro, 3',5'-dimethyl and 3'-methyl-4'-fluoro.

Preferred examples for R⁵ include primary and secondary C₁₋₆ alkyl and C₃₋₆ cycloalkyl.

Preferred examples for Y include -CH₂-CH₂- and -CH=CH-.

Preferred examples for Z include



-CH(OH)CH₂CH₂(OH)CH₂CO₂R¹², -CH(OH)CH₂C(O)CH₂CO₂R¹² and -CH(OH)CH₂C(OR¹³)₂CH₂CO₂R¹².

Now, more preferred substituents of the compounds of the present invention will be described.

As more preferred examples for R¹, R² and R⁶, when both R² and R⁶ are hydrogen, R¹ is hydrogen, 5-fluoro, 6-fluoro, 7-fluoro, 8-fluoro, 5-chloro, 6-chloro, 7-chloro, 8-chloro, 5-bromo, 6-bromo, 7-bromo, 8-bromo, 5-methyl, 6-methyl, 7-methyl, 8-methyl, 5-methoxy, 6-methoxy, 7-methoxy, 8-methoxy, 5-trifluoromethyl, 6-trifluoromethyl, 7-trifluoromethyl, 8-trifluoromethyl, 6-trifluoromethoxy, 6-difluoromethoxy, 8-hydroxyethyl, 5-hydroxy, 6-hydroxy, 7-hydroxy, 8-hydroxy, 6-ethyl, 6-n-butyl and 7-dimethylamino.

When R⁶ is hydrogen, R¹ and R² together represent 6-chloro-8-methyl, 6-bromo-7-methoxy, 6-methyl-7-chloro, 6-chloro-8-hydroxy, 5-methyl-2-hydroxy, 6-methoxy-7-chloro, 6-chloro-7-methoxy, 6-hydroxy-7-chloro, 6-chloro-7-hydroxy, 6-chloro-8-bromo, 5-chloro-6-hydroxy, 6-bromo-8-chloro, 6-bromo-8-hydroxy, 5-methyl-8-chloro, 7-hydroxy-8-chloro, 6-bromo-8-hydroxy, 6-methoxy-7-methyl, 6-chloro-8-bromo, 6-methyl-8-bromo, 6,7-difluoro, 6,8-difluoro, 6,7-methylenedioxy, 6,8-dichloro, 5,8-dimethyl, 6,8-dimethyl, 6,7-dimethoxy, 6,7-diethoxy, 6,7-dibromo or 6,8-dibromo.

When R¹, R² and R⁶ are not hydrogen, they together represent 5,7-dimethoxy-8-hydroxy, 5,8-dichloro-6-hydroxy, 6,7,8-trimethoxy, 6,7,8-trimethyl, 6,7,8-trichloro, 5-fluoro-6,8-dibromo or 5-chloro-6,8-dibromo.

As more preferred examples for R³ and R⁴, when R³ is hydrogen, R⁴ is hydrogen, 4'-methyl, 4'-chloro or 4'-fluoro. When both R³ and R⁴ are not hydrogen, they together represent 3',5'-dimethyl or 3'-methyl-4'-fluoro.

As more preferred examples for R⁵, the above-mentioned preferred examples of R⁵ may be mentioned.

As preferred examples for Y, -CH₂-CH₂- and (E)-CH=CH- may be mentioned. As more preferred examples for Z, the above preferred examples for Z may be mentioned.

Now, still further preferred substituents of the compounds of the present invention will be described. As examples for R¹, R² and R⁶, when both R² and R⁶ are hydrogen, R¹ is hydrogen, 6-methyl, 6-ethyl, 6-trifluoromethyl, 6-hydroxy, 6-methoxy, 6-chloro, 6-bromo, 6-n-butyl and 7-dimethylamino.

When only R⁶ is hydrogen, R¹ and R² represent 6,8-dichloro, 5,8-dimethyl, 6,8-dimethyl, 6,7-dimethoxy, 6,7-diethoxy, 6,7-dibromo, 6,8-dibromo, 6,7-difluoro and 6,8-difluoro.

As still further preferred examples for R³ and R⁴, when R³ is hydrogen, R⁴ is hydrogen, 4'-chloro or 4'-fluoro, or R³ and R⁴ together represent 3'-methyl-4'-fluoro.

Still further preferred examples for R⁵ include ethyl, n-propyl, i-propyl and cyclopropyl.

Still further preferred examples for Y include (E)-CH=CH-.

As still further preferred examples for Z, the above-mentioned preferred example for Z may be mentioned.

Now, the most preferred substituents for the compounds of the present invention will be described.

As the most preferred examples for R¹, R² and R⁶, when both R² and R⁶ are hydrogen, R¹ is hydrogen, 6-methyl or 6-chloro.

When only R⁶ is hydrogen, R¹ and R² together represent, for example, 6,7-dimethoxy.

As the most preferred examples for R³ and R⁴, R³ is hydrogen and R⁴ is hydrogen, 4'-chloro or 4'-fluoro.

The most preferred examples for R⁵ include i-propyl and cyclopropyl. The most preferred example for Y may be (E)-CH=CH-.


As the most preferred examples for Z, the above-mentioned preferred examples for Z may be mentioned.


Now, particularly preferred specific compounds of the present invention will be presented. The following compounds (a) to (z) are shown in the form of carboxylic acids. However, the present invention include not only the compounds in the form of carboxylic acids but also the corresponding lactones formed by the condensation of the carboxylic acids with hydroxy at the 5-position, and sodium salts and lower alkyl esters (such as methyl, ethyl, i-propyl and n-propyl esters) of the carboxylic acids, which can be physiologically hydrolyzed to the carboxylic acids.

(a) (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-(1''-methylene)-quinolin-3'-yl]-hept-6-enoic acid

- (b) (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-(1''-methylethyl)-6'-chloro-quinolin-3'-yl]-hept-6-enoic acid
- (c) (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-(1''-methylethyl)-6'-methyl-quinolin-3'-yl]-hept-6-enoic acid
- 5 (d) (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-(1''-methylethyl)-6',7'-dimethoxy-quinolin-3'-yl]-hept-6-enoic acid
- (e) (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-cyclopropyl-quinolin-3'-yl]-hept-6-enoic acid
- (f) (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-cyclopropyl-6'-chloro-quinolin-3'-yl]-hept-6-enoic acid
- (g) (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-cyclopropyl-6'-methyl-quinolin-3'-yl]-hept-6-enoic acid
- 10 (h) (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-cyclopropyl-6',7'-dimethoxy-quinolin-3'-yl]-hept-6-enoic acid
- (i) (E)-3,5-dihydroxy-7-[4'-(4''-chlorophenyl)-2'-(1''-methylethyl)-quinolin-3'-yl]-hept-6-enoic acid
- (j) (E)-3,5-dihydroxy-7-[4'-(4''-chlorophenyl)-2'-(1''-methylethyl)-6'-chloro-quinolin-3'-yl]-hept-6-enoic acid
- 15 (k) (E)-3,5-dihydroxy-7-[4'-(4''-chlorophenyl)-2'-(1''-methylethyl)-6'-methyl-quinolin-3'-yl]-hept-6-enoic acid
- (l) (E)-3,5-dihydroxy-7-[4'-(4''-chlorophenyl)-2'-(1''-methylethyl)-6',7'-dimethoxy-quinolin-3'-yl]-hept-6-enoic acid
- (m) (E)-3,5-dihydroxy-7-[4'-(4''-chlorophenyl)-2'-cyclopropyl-quinolin-3'-yl]-hept-6-enoic acid
- 20 (n) (E)-3,5-dihydroxy-7-[4'-(4''-chlorophenyl)-2'-cyclopropyl-6'-chloro-quinolin-3'-yl]-hept-6-enoic acid
- (o) (E)-3,5-dihydroxy-7-[4'-(4''-chlorophenyl)-2'-cyclopropyl-6'-methyl-quinolin-3'-yl]-hept-6-enoic acid
- (p) (E)-3,5-dihydroxy-7-[4'-(4''-chlorophenyl)-2'-cyclopropyl-6',7'-dimethoxy-quinolin-3'-yl]-hept-6-enoic acid
- (q) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-(1''-methylethyl)-quinolin-3'-yl]-hept-6-enoic acid
- 25 (r) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-(1''-methylethyl)-6'-chloro-quinolin-3'-yl]-hept-6-enoic acid
- (s) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-(1''-methylethyl)-6'-methyl-quinolin-3'-yl]-hept-6-enoic acid
- (t) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-(1''-methylethyl)-6',7'-dimethoxy-quinolin-3'-yl]-hept-6-enoic acid
- (u) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-cyclopropyl-quinolin-3'-yl]-hept-6-enoic acid
- (v) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-cyclopropyl-6'-chloro-quinolin-3'-yl]-hept-6-enoic acid
- 30 (w) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-cyclopropyl-6'-methyl-quinolin-3'-yl]-hept-6-enoic acid
- (x) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-cyclopropyl-6',7'-dimethoxy-quinolin-3'-yl]-hept-6-enoic acid
- (y) (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-(1''-methylethyl)-6'-methoxy-quinolin-3'-yl]-hept-6-enoic acid
- 35 (z) (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-cyclopropyl-6'-methoxy-quinolin-3'-yl]-hept-6-enoic acid

The mevalonolactones of the formula I can be prepared by the following reaction scheme. The enal III can also be prepared by processes K, L and M.





In the above reaction scheme, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^{12} are as defined above with respect to the formula I, and R^{21} and R^{22} independently represent C_{1-4} lower alkyl such as methyl, ethyl, *n*-propyl, *i*-propyl or *n*-butyl.

Step A represents a reduction reaction of the ester to a primary alcohol. Such reduction reaction can be conducted by using various metal hydrides, preferably diisobutylaluminum hydride, in a solvent such as tetrahydrofuran or toluene at a temperature of from -20 to 20°C , preferably from -10 to 10°C .

Step B represents an oxidation reaction of the primary alcohol to an aldehyde, which can be conducted by using various oxidizing agents. Preferably, the reaction can be conducted by using pyridinium chlorochromate in methylene chloride at a temperature of from 0 to 25°C , or by using oxalyl chloride, dimethyl sulfoxide and a tertiary amine such as triethylamine (Swern oxidation), or by using a sulfur trioxide

pyridine complex.

Step C represents a synthesis of a 3-ethoxy-1-hydroxy-2-propene derivative, which can be prepared by reacting a compound V to lithium compound which has been preliminarily formed by treating cis-1-ethoxy-2-(tri-n-butylstannyl)ethylene with butyl lithium in tetrahydrofuran.

5 As the reaction temperature, it is preferred to employ a low temperature at a level of from -60 to -78 °C.

Step D represents a synthesis of an enal by acidic hydrolysis. As the acid catalyst, it is preferred to employ p-toluene sulfonic acid, hydrochloric acid or sulfuric acid, and the reaction may be conducted in a solvent mixture of water and tetrahydrofuran or ethanol at a temperature of from 10 to 25 °C. The 3-ethoxy-1-hydroxy-2-propene derivative obtained in Step C can be used in Step D without purification i.e. by simply removing tetra-n-butyl tin formed simultaneously.

Step E represents a double anion condensation reaction between the enal III and an acetoacetate. Such condensation reaction is preferably conducted by using sodium hydride and n-butyl lithium as the base in tetrahydrofuran at a temperature of from -80 to 0 °C, preferably from -30 to -10 °C.

Step F represents a reduction reaction of the carbonyl group, which can be conducted by using a metal hydride, preferably sodium borohydride in ethanol at a temperature of from -10 to 25 °C, preferably from -10 to 5 °C.

Further, the reduction reaction may be conducted by using zinc borohydride in dry ethyl ether or dry tetrahydrofuran at a temperature of -100 to 25 °C, preferably from -80 to -50 °C.

Step G is a step for hydrolyzing the ester. The hydrolysis can be conducted by using an equimolar amount of a base, preferably potassium hydroxide or sodium hydroxide, in a solvent mixture of water and methanol or ethanol at a temperature of from 10 to 25 °C. The free acid hereby obtained may be converted to a salt with a suitable base.

Step H is a step for forming a mevalonolactone by the dehydration reaction of the free hydroxy acid I-2. The dehydration reaction can be conducted in benzene or toluene under reflux while removing the resulting water or by adding a suitable dehydrating agent such as molecular sieve.

Further, the dehydration reaction may be conducted in dry methylene chloride by using a lactone-forming agent such as carbodiimide, preferably a water soluble carbodiimide such as N-cyclohexyl-N'-[2'-(methylmorpholinium)ethyl]carbodiimide p-toluene sulfonate at a temperature of from 10 to 35 °C, preferably from 20 to 25 °C.

Step J represents a reaction for hydrogenating the double bond connecting the mevalonolactone moiety and the quinoline ring. This hydrogenation reaction can be conducted by using a catalytic amount of palladium-carbon or rhodium-carbon in a solvent such as methanol, ethanol, tetrahydrofuran or acetonitrile at a temperature of from 0 to 50 °C, preferably from 10 to 25 °C.

Step K represents a reaction for the synthesis of an α,β -unsaturated carboxylic acid ester, whereby a trans-form α,β -unsaturated carboxylic acid ester can be obtained by a so-called Horner-Wittig reaction by using an alkoxycarbonylmethyl phosphonate. The reaction is conducted by using sodium hydride or potassium t-butoxide as the base in dry tetrahydrofuran at a temperature of from -30 to 0 °C, preferably from -20 to -15 °C.

Step L represents a reduction reaction of the α,β -unsaturated carboxylic acid ester to an allyl alcohol. This reduction reaction can be conducted by using various metal hydrides, preferably diisobutylaluminumhydride, in a solvent such as dry tetrahydrofuran or toluene at a temperature of from -10 to 10 °C, preferably from -10 to 0 °C.


Step M represents an oxidation reaction of the allyl alcohol to an enal. This oxidation reaction can be conducted by using various oxidizing agents, particularly active manganese dioxide, in a solvent such as tetrahydrofuran, acetone, ethyl ether or ethyl acetate at a temperature of from 0 to 100 °C, preferably from 15 to 50 °C.

Step N represents a reaction for the synthesis of an α,β -unsaturated ketone by the selective oxidation of the dihydroxy carboxylic acid ester. This reaction can be conducted by using activated manganese dioxide in a solvent such as ethyl ether, tetrahydrofuran, benzene or toluene at a temperature of from 20 to 80 °C, preferably from 40 to 80 °C.

In addition to the compounds disclosed in Examples given hereinafter, compounds of the formulas I-2 and I-5 given in Table 1 can be prepared by the process of the present invention. In Table 1, i- means iso, sec- means secondary and c- means cyclo. Likewise, Me means methyl, Et means ethyl, Pr means propyl, Bu means butyl, Pent means pentyl, Hex means hexyl and Ph means phenyl.

55

Table 1



R^1	R^2	R^3	R^4	R^5	R^6
6-OMe	H	H	H	i-Pr	H
6-OMe	H	4-F	H	i-Pr	H
6-Br	H	4-F	H	i-Pr	H
6-Me	8-Me	4-F	H	i-Pr	H
7-OMe	8-OMe	4-F	H	i-Pr	H
6-Br	H	2-F	H	i-Pr	H
6,7					
H	H	4-F	H	i-Pr	H
35					
H	H	4-F	H		H
H	H	4-Ph	H	i-Pr	H
H	H	4-PhCH ₂	H	i-Pr	H
6-Cl	H	4-F	H	c-Pr	H
6-Cl	H	4-F	H	sec-Bu	H
6-OCH ₂ Ph	H	4-F	H	i-Pr	H
H	H	4-F	H	i-Bu	H
H	H	4-F	H	c-Pent	H
6-Cl	H	4-F	H	c-Pent	H
6-Me ₂ N	H	4-F	H	i-Pr	H

	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶
5	6-Me	H	4-F	H	c-Pr	H
	6-i-Pr	H	4-F	H	i-Pr	H
10	7-Me	H	4-F	H	c-Pr	H
	6-OMe	H	4-F	H	c-Pr	H
	6-Br	H	4-F	H	c-Pr	H
15	6-i-Pr	H	4-F	H	c-Pr	H
	6-C ℓ	8-C ℓ	4-F	H	c-Pr	H
20	5-F	6-Br	4-F	H	i-Pr	8-Br
	6-OMe	7-OMe	4-F	H	i-Pr	8-OMe
25	6-Me	7-Me	4-F	H	i-Pr	8-Me
	6-C ℓ	7-C ℓ	4-F	H	i-Pr	8-C ℓ
	H	H	4-F	H	c-Bu	H
30	H	H	4-F	H	c-Hex	H
	6-OMe	7-OMe	H	H	i-Pr	H
35	6-OMe	7-OMe	4-C ℓ	H	i-Pr	H
	6-OMe	7-OMe	H	H	c-Pr	H
40	6-OMe	7-OMe	4-C ℓ	H	c-Pr	H
	6-OMe	7-OMe	4-F	H	c-Pr	H

45

50

55

	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶
5	6-Me	H	H	H	i-Pr	H
	6-Me	H	4-Cl	H	i-Pr	H
	6-Me	H	H	H	c-Pr	H
10	6-Me	H	4-Cl	H	c-Pr	H
	6-Me	H	4-F	H	c-Pr	H
15	6-Cl	H	H	H	i-Pr	H
	6-Cl	H	4-Cl	H	i-Pr	H
	6-Cl	H	H	H	c-Pr	H
20	6-Cl	H	4-Cl	H	c-Pr	H
	6-Cl	H	4-F	H	c-Pr	H
25	H	H	H	H	i-Pr	H
	H	H	4-Cl	H	i-Pr	H
30	H	H	H	H	c-Pr	H
	H	H	4-Cl	H	c-Pr	H
35	H	H	4-F	H	c-Pr	H

Further, pharmaceutically acceptable salts such as potassium salts or esters such as ethyl esters or methyl esters of these compounds can be prepared in the same manner.

The compounds of the present invention exhibit high inhibitory activities against the cholesterol biosynthesis wherein HMG-CoA reductase acts as a rate-limiting enzyme, as shown by the test results given hereinafter, and thus are capable of suppressing or reducing the amount of cholesterol in blood as lipoprotein. Thus, the compounds of the present invention are useful as curing agents against hyperlipidemia, hyperlipoproteinemia and atherosclerosis.

They may be formulated into various suitable formulations depending upon the manner of the administration. The compounds of the present invention may be administered in the form of free acids or in the form of physiologically hydrolyzable and acceptable esters or lactones, or pharmaceutically acceptable salts.

The pharmaceutical composition of the present invention is preferably administered orally in the form of the compound of the present invention per se or in the form of powders, granules, tablets or capsules formulated by mixing the compound of the present invention with a suitable pharmaceutically acceptable carrier including a binder such as hydroxypropyl cellulose, syrup, gum arabic, gelatin, sorbitol, tragacanth gum, polyvinyl pyrrolidone or CMC-Ca, an excipient such as lactose, sugar, corn starch, calcium phosphate, sorbitol, glycine or crystal cellulose powder, a lubricant such as magnesium stearate, talk, polyethylene glycol or silica, and a disintegrator such as potato starch.

However, the pharmaceutical composition of the present invention is not limited to such oral administration and it is applicable for parenteral administration. For example, it may be administered in the form of e.g. a suppository formulated by using oily base material such as cacao butter, polyethylene glycol, lanolin or fatty acid triglyceride, a transdermal therapeutic base formulated by using liquid paraffin, white vaseline,

a higher alcohol, Macrogol ointment, hydrophilic ointment or hydro-gel base material, an injection formulation formulated by using one or more materials selected from the group consisting of polyethylene glycol, hydro-gel base material, distilled water, distilled water for injection and excipient such as lactose or corn starch, or a formulation for administration through mucous membranes such as an ocular mucous membrane, a nasal mucous membrane and an oral mucous membrane.

Further, the compounds of the present invention may be combined with basic ion-exchange resins which are capable of binding bile acids and yet not being absorbed in gastrointestinal tract.

The daily dose of the compound of the formula I is from 0.05 to 500 mg, preferably from 0.5 to 50 mg for an adult. It is administered from once to three times per day. The dose may of course be varied depending upon the age, the weight or the condition of illness of the patient.

The compounds of the formulas II to VII are novel, and they are important intermediates for the preparation of the compounds of the formula I. Accordingly, the present invention relates also to the compounds of the formulas II to VII and the processes for their production.

Now, the present invention will be described in further detail with reference to Test Examples for the pharmacological activities of the compounds of the present invention, their Preparation Examples and Formulation Examples. However, it should be understood that the present invention is by no means restricted by such specific Examples.

20 PHARMACOLOGICAL TEST EXAMPLES

25 Test A: Inhibition of cholesterol biosynthesis from acetate in vitro

Enzyme solution was prepared from liver of male Wistar rat bilially cannulated and discharged bile for over 24 hours. Liver was cut out at mid-dark and microsome and supernatant fraction which was precipitable with 40-80% of saturation of ammonium sulfate (sup fraction) were prepared from liver homogenate according to the modified method of Knauss et. al.; Kuroda, M., et. al., Biochim. Biophys. Acta, 489, 119 (1977). For assay of cholesterol biosynthesis, microsome (0.1 mg protein) and sup fraction (1.0 mg protein) were incubated for 2 hours at 37°C in 200 µl of the reaction mixture containing ATP; 1 mM, Glutathione; 6 mM, Glucose-1-phosphate; 10 mM, NAD; 0.25 mM, NADP; 0.25 mM, CoA; 0.04 mM and 0.2 mM [2-¹⁴C]sodium acetate (0.2 µCi) with 4 µl of test compound solution dissolved in water or dimethyl sulfoxide. To stop reaction and saponify, 1 ml of 15% EtOH-KOH was added to the reactions and heated at 35 75°C for 1 hour. Nonsaponifiable lipids were extracted with petroleum ether and incorporated ¹⁴C radioactivity was counted. Inhibitory activity of compounds was indicated with IC50.

40 Test B: Inhibition of cholesterol biosynthesis in culture cells

Hep G2 cells at over 5th passage were seeded to 12 well plates and incubated with Dulbecco's modified Eagle (DME) medium containing 10% of fetal bovine serum (FBS) at 37°C, 5% CO₂ until cells were confluent for about 7 days. Cells were exposed to the DME medium containing 5% of lipoprotein deficient serum (LPDS) prepared by ultracentrifugation method for over 24 hours. Medium was changed to 45 0.5 ml of fresh 5% LPDS containing DME before assay and 10 µl of test compound solution dissolved in water or DMSO were added. 0.2 µCi of [2-¹⁴C]sodium acetate (20 µl) was added at 0 hr(B-1) or 4 hrs(B-2) after addition of compounds. After 4 hrs further incubation with [2-¹⁴C]sodium acetate, medium was removed and cells were washed with phosphate buffered saline(PBS) chilled at 4°C. Cells were scraped with rubber policeman and collected to tubes with PBS and digested with 0.2 ml of 0.5 N KOH at 37°C. 50 Aliquot of digestion was used for protein analysis and remaining was saponified with 1 ml of 15% EtOH-KOH at 75°C for 1 hour. Nonsaponifiable lipids were extracted with petroleum ether and ¹⁴C radioactivity was counted. Counts were revised by cell protein and indicated with DPM/mg protein. Inhibitory activity of compounds was indicated with IC50.

55

Test C: Inhibition of cholesterol biosynthesis in vivo

Male Sprague-Dawley rats weighing about 150 g were fed normal Purina chow diet and water ad libitum, and exposed to 12 hours light/12 hours dark lighting pattern (2:00 PM - 2:00 AM dark) prior to use for in vivo inhibition test of cholesterol biosynthesis. Animals were separated groups consisting of five rats as to be average mean body weight in each groups. Test compounds at dosage of 0.02-0.2 mg/kg body weight (0.4 ml/100 g body weight), were dissolved in water or suspended or in 0.5% methyl cellulose and orally administered at 2-3 hours before mid-dark (8:00 PM), while cholesterol biosynthesis reaches to maximum in rats. As control, rats were orally administered only water or vehicle. At 90 minutes after sample administration, rats were injected intraperitoneally with 10 μ Ci of [2- 14 C]sodium acetate at volume of 0.2 ml per one. 2 Hours later, blood samples were obtained and serum were separated immediately. Total lipids were extracted according to the method of Folch et al. and saponified with EtOH-KOH. Nonsaponifiable lipids were extracted with petroleum ether and radio activity incorporated into nonsaponifiable lipids was counted.

Inhibitory activity was indicated as percent decrease of counts in testing groups (DPM/2 ml serum/2 hours) from that in control group.

With respect to the compounds of the present invention, the inhibitory activities against the cholesterol biosynthesis in which HMG-CoA reductase serves as a rate limiting enzyme, were measured by the above Test A and B. The results are shown in Tables, 2, 2-2, 3 and 3-2. Further, the results of the measurements by Test C are also presented.

Table 2

Inhibitory activities by Test A	
Compound	I ₅₀ (molar concentration)
(Compounds of the present invention)	
I-13	1.25×10^{-7}
I-51	1.0×10^{-8}
I-52	7.1×10^{-8}
I-53	1.9×10^{-7}
(Reference compounds)	
Mevinolin	1.4×10^{-8}
CS-514	9.0×10^{-9}


In Table 2-2, the relative activities are shown based on the activities of CS-514 being evaluated to be 1.

Table 2-2

Relative activities by Test A	
Compound	Relative activities
(Compounds of the present invention)	
I-16	1.75
I-116	2.25
I-117	0.37
I-120	3.21
I-522	0.76

Structures of reference compounds:

(1) Mevinolin



(2) CS-514




Table 3

Inhibitory activities by Test B-1	
Compound	I ₅₀ (molar concentration)
(Compound of the present invention)	
I-51	1 x 10 ⁻⁷
(Reference compound)	
CS-514	3.5 x 10 ⁻⁷

In Table 3-2, the relative activities are shown based on the activities of CS-514 being evaluated to be 1.

Table 3-2

Relative activities by Test B-1	
Compound	Relative activities
I-116	19.4
I-520	20.0
II-20	20.8

Results of the measurement of the inhibitory activities by Test C

The percent decrease of counts after the oral administration of 0.05 mg/kg of compound I-520 was 55% relative to the measured value of the control group. The percent decrease of counts after the oral administration of 10 mg/kg of CS-514 was 55% under the same condition. The compounds of the present invention exhibited activities superior to the reference compound such as CS-514 or Mevinolin in Test A, and exhibited activities superior to CS-514 in Tests B and C.

Test D: Acute toxicity

A 0.5% CMC suspension of a test compound was orally administered to ICR male mice (group of three mice). The acute toxicity was determined based on the mortality after seven days. With compound I-57, I-58, I-59, I-511, I-512, I-513, I-514, I-515, I-517 and I-523 of the present invention, the mortality was 0% even when they were orally administered in an amount of 1000 mg/kg.

EXAMPLE 1

Ethyl (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-(1'''-methylethyl)-quinolin-3'-yl]-hept-6-enoate (compound I-11) (prepared by steps of Example 1-a through Example 1-q)

EXAMPLE 1-a: Ethyl 4-(4'-fluorophenyl)-2-(1'-methylethyl)-quinolin-3-yl-carboxylate (compound VII-1)

The synthesis was conducted in accordance with the method disclosed in J. Org. Chem., 2899 (1966). 6.45 g (0.03 mol) of 2-amino-4'-fluorobenzophenone, 5.53 g (0.035 mol) of ethyl isobutyrylacetate and 0.1 ml of conc. sulfuric acid were dissolved in 30 ml of glacial acetic acid, and the mixture was heated at 100°C for about 10 hours. After confirming the substantial disappearance of 2-amino-4'-fluorobenzophenone by thin layer chromatography, the reaction solution was cooled to room temperature, and a mixture of 45 ml of conc. aqueous ammonia and 120 ml of water cooled with ice, was gradually added thereto. A separated oily substance was solidified when left to stand overnight in a refrigerator. This solid was recrystallized from a small amount of ethanol to obtain 6.47 g (55%) of white powder. Melting point: 68-70.5°C

EXAMPLE 1-b: 4-(4'-fluorophenyl)-3-hydroxymethyl-2-(1'-methylethyl)-quinoline (compound VI-1)

5.4 g (0.016 mol) of compound VII-1 was dissolved in dry toluene under a nitrogen atmosphere and cooled in ice bath to 0°C. To this solution, 40 ml of a 16 wt% diisobutylaluminum hydride-toluene solution was dropwis added, and the mixture was stirred at 0°C for two hours. After confirming the complete

disappearance of compound VII-1 by thin layer chromatography, a saturated ammonium chloride solution was added thereto at 0 °C to terminate the reaction. Ethyl ether was added to the reaction mixture, and the organic layer was separated. A gelled product was dissolved by an addition of an aqueous sodium hydroxide solution and extracted anew with ethyl ether. The ethyl ether extracts were put together, dried
 5 over anhydrous magnesium sulfate and filtered. The solvent was distilled off. The residual oil underwent crystallization when left to stand. It was recrystallized from ethyl acetate-n-hexane to obtain 3.3 g of white crystals. Yield: 70%. Melting point: 136-137 °C.

10 EXAMPLE 1-c: 4-(4'-fluorophenyl)-2-(1'-methylethyl)-quinolin-3-yl-carboxyaldehyde (compound V-1)

2.0 g (9.3 mmol) of pyridinium chlorochromate and 0.4 g of anhydrous sodium acetate was suspended in 10 ml of dry dichloromethane. To this suspension, a solution obtained by dissolving 1 g (3.4 mmol) of compound VI-1 in 10 ml of dry dichloromethane, was immediately added at room temperature. The mixture
 15 was stirred for one hour. Then, 100 ml of ethyl ether was added thereto, and the mixture was thoroughly mixed. The reaction mixture was filtered under suction through a silica gel layer. The filtrate was dried under reduced pressure. The residue was dissolved in the isopropyl ether, and insoluble substances were filtered off. The filtrate was again dried under reduced pressure, and the residue was recrystallized from diisopropyl ether to obtain 0.7 g (Yield: 70%) of slightly yellow prism crystals. Melting point: 124-126 °C.

20 EXAMPLE 1-d: 3-(3'-ethoxy-1'-hydroxy-2'-propenyl)-4-(4'-fluorophenyl)-2-(1'-methylethyl)-quinoline (compound IV-1)

25 1.13 g (3.13 mmol) of cis-1-ethoxy-2-(tri-n-butylstannyl)ethylene was dissolved in 8 ml of dry tetrahydrofuran, and the solution was cooled to -78 °C in a nitrogen stream. To this solution, 2 ml (3.2 mmol) of a 15 wt% n-butyllithium-n-hexane solution was dropwise added. The mixture was stirred for 45 minutes. Then, a solution prepared by dissolving 0.76 g (2.6 mmol) of compound V-1 in 10 ml of dry tetrahydrofuran was dropwise added thereto. The reaction mixture was stirred at -78 °C for two hours. Then,
 30 2 ml of a saturated ammonium chloride solution was added thereto to terminate the reaction. The organic layer was extracted with diethyl ether, and the diethyl ether extract was washed with a saturated sodium chloride aqueous solution and dried over anhydrous magnesium sulfate. The solvent was distilled off under reduced pressure. The residue was separated with n-hexane and acetonitrile. The solvent was distilled off under reduced pressure from the acetonitrile layer, and an oily substance thereby obtained was purified by
 35 silica gel column chromatography (eluent: 2.5% methanol-chloroform) to obtain 0.91 g of the desired compound in a purified oily form.

H-MNR (CDCl₃) δ ppm:

1.1(t,3H,7Hz) 1.37(d,6H,J=7Hz) 3.7(m,1H) 3.7(q,2H,J=7Hz) 4.75(t,1H,7Hz) 5.7(m,1H) 5.95(m,1H) 7.05-8.2-
 (m,8H)

40 EXAMPLE 1-e: (E)-3-[4'-(4''-fluorophenyl)-2'-(1''-methylethyl)-quinolin-3'-yl]propenaldehyde (compound III-1)

45 0.91 g of compound IV-1 was dissolved in 20 ml of tetrahydrofuran, and 5 ml of water and 100 mg of p-toluenesulfonic acid were added thereto. The mixture was stirred at room temperature for 24 hours. The reaction solution was extracted with diethyl ether a few times. The extracts were washed with a saturated sodium chloride aqueous solution and dried over anhydrous magnesium sulfate. Then, the solvent was distilled off. The residue was purified by silica gel column chromatography (eluent: chloroform) to obtain the
 50 desired product as white prism crystals. 0.4 g (50%). Melting point: 127-128 °C.

EXAMPLE 1-f: Ethyl (E)-7-[4'-(4''-fluorophenyl)-2'-(1''-methylethyl)-quinolin-3'-yl]-5-hydroxy-3-oxohepto-6-enoate (compound II-1)

55 50 mg of 60% sodium hydride was washed with dry petroleum ether and dried under a nitrogen stream, and then suspended in 5 ml of dry tetrahydrofuran. The suspension was cooled to -15 °C in a nitrogen atmosphere. Then, 120 mg (0.92 mmol) of ethyl acetoacetate was dropwise added thereto, and the mixture

was stirred for 15 minutes. Then, 0.6 ml (0.92 mmol) of a 15 wt% n-butyllithium-n-hexane solution was dropwise added thereto, and the mixture was stirred for 30 minutes. Then, a solution prepared by dissolving 160 mg (0.5 mmol) of compound II-1 in dry tetrahydrofuran, was dropwise added thereto, and the mixture was stirred for one hour. To the reaction mixture, 1 ml of a saturated ammonium chloride aqueous solution was added at -15 °C. Then, the mixture was extracted three times with diethyl ether. The diethyl ether solution was washed with a saturated sodium chloride aqueous solution and dried over anhydrous magnesium sulfate. The solution was evaporated to dryness under reduced pressure. The residue was recrystallized from diisopropyl ether to obtain 130 mg (yield: 59%) of white crystals. Melting point: 99-101 °C.

10

EXAMPLE 1-g: Ethyl (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-(1''-methylethyl)-quinolin-3'-yl]-hept-6-enoate (compound I-11)

110 mg (0.245 mmol) of compound II-1 was dissolved in 5 ml of ethanol in a nitrogen atmosphere, and the solution was cooled 0 °C. Then, 10 mg (0.263 mmol) of sodium borohydride was added, and the mixture was stirred for one hour. Then, 1 ml of a 10% hydrochloric acid aqueous solution was added thereto, and the mixture was extracted three times with ethyl ether. The ethyl ether solution was washed with a saturated sodium chloride aqueous solution and dried over anhydrous magnesium sulfate. Then, the solution was evaporated to dryness under reduced pressure. The residual oil was purified by silica gel column chromatography (eluent: 5% methanol-chloroform) to obtain the desired product as a pure colorless oily substance. 70 mg (Yield: 64%)

H-NMR (CDCl₃) δ ppm:

1.30(t,3H,J=8Hz) 1.39(d,6H,J=8Hz) 1.4-1.8(m,2H) 2.42(d,2H,J=7Hz) 3.0-3.8 (m,2H) 3.50(m,1H) 3.9-4.6-(m,2H) 4.20(q,2H,J=8Hz) 5.35(m,1H) 6.59(m,1H) 7.10-8.18(m,8H)

25

EXAMPLE 2

30

Sodium salt of (E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-(1''-methylethyl)-quinolin-3'-yl]-hept-6-enoic acid (compound I-51)

60 mg (0.133 mmol) of compound I-11 was dissolved in 3 ml of ethanol. Then, 0.26 ml of a 0.5 N sodium hydroxide aqueous solution was dropwise added thereto. The mixture was stirred at room temperature for further one hour, and ethanol was distilled off under reduced pressure. Then, 5 ml of water was added thereto, and the mixture was extracted with ethyl ether. The aqueous layer was freeze-dried to obtain 40 mg (67%) of hygroscopic white powder. Melting point: 207-209 °C (decomposed).

40

EXAMPLE 3

45

(E)-3,5-dihydroxy-7-[4'-(4''-fluorophenyl)-2'-(1''-methylethyl)-quinolin-3'-yl]-hept-6-enoic acid (compound I-21)

110 mg (0.244 mmol) of compound I-11 was dissolved in 10 ml of ethanol. Then, 0.79 ml of a 0.5 N sodium hydroxide aqueous solution was dropwise added thereto. The mixture was stirred at room temperature for further one hour, and ethanol was distilled off under reduced pressure. Then, 10 ml of water was added thereto, and the mixture was extracted with ethyl ether. The aqueous layer was weakly acidified (pH 4) with a dilute hydrochloric aqueous solution and extracted three times with ethyl ether. The ethyl ether layers were put together and dried over anhydrous magnesium sulfate. Then, the solvent was distilled off under reduced pressure to obtain 90 mg of slightly yellow oily substance.

55

H-NMR (CDCl₃) δ ppm:

1.36(d,6H,J=7Hz) 2.4(m,2H) 3.5(m,1H) 3.45(m,1H) 3.8-4.6(m,2H) 5.40(dd,1H,J₁=19Hz,J₂=8Hz) 6.55(d,1H,J=19Hz) 7.0-8.3(m,8H)

EXAMPLE 4

5 (E)-6-[4'-(4''-fluorophenyl)-2'-(1''-methylethyl)-quinolin-3'-ylethenyl]-4-hydroxy-3,4,5,6-tetrahydro-2H-pyran-2-one (compound I-31)

90 mg of compound I-21 was dissolved in 10 ml of dry toluene, and the solution was refluxed under heating for 3 hours by means of a Dean Stark apparatus.

10 Toluene was distilled off under reduced pressure, and the residual solid was recrystallized from diisopropyl ether to obtain 40 mg of colorless prism crystals. Melting point: 182-184°C.

By silica gel thin chromatography, the product gave two absorption spots close to each other attributable to the diastereomers. (Developing solvent: 3% methanol-chloroform)

15 These diastereomers were separated and isolated by silica gel thin layer chromatography. [Developing solvent: t-BuOMe/hexane/acetone = 7/2/1 (v/v), R_f = 0.6 and 0.7 (obtained weight ratio: 1/2)]

R_f = 0.7: trans lactone

20 H-NMR (CDCl₃) δ ppm:

1.40(d,6H,J = 7Hz) 1.6(m,2H) 2.65(m,2H) 3.48(m,1H) 4.20(m,1H) 5.15(m,1H) 5.37(dd,1H,J₁ = 18Hz,J₂ = 7Hz) 6.68(d,1H,J = 19Hz) 7.1-8.2(m,8H)

25

R_f = 0.6: cis lactone

H-NMR (CDCl₃) δ ppm:

30 1.40(d,6H,J = 7Hz) 1.6(m,2H) 2.65(m,2H) 3.48(m,1H) 4.20(m,1H) 4.65(m,1H) 5.40(dd,1H,J₁ = 18Hz,J₂ = 7Hz) 6.66(m,1H) 7.0-8.2(m,8H)

EXAMPLE 5

35

6-[4'-(4''-fluorophenyl)-2'-(1''-methylethyl)-quinolin-3'-ylethynyl]-4-hydroxy-3,4,5,6-tetrahydro-2H-pyran-2-one (compound I-41)

40

20 mg of a mixture of diastereomers of compound I-31 was dissolved in 5 ml of ethanol, and 10 mg of 5% palladium-carbon was added thereto. The mixture was stirred under a hydrogen atmosphere. After confirming the disappearance of the starting substance and the appearance of a new spot by thin layer chromatography, the palladium-carbon was filtered off, and ethanol was distilled off to obtain colorless oil.

45 This oil was purified by preparative thin layer chromatography to obtain 16 mg of the desired product as pure colorless oil.

MS(m/e): 408(M⁺ + H), 407(M⁺), 366, 292, 278

In the same manner as in Example 1-a, compounds VII-2 to VII-27 were prepared. The physical properties of these compounds are shown in Table 4. (In the Table, R¹, R², R³, R⁴, R⁵ and R²¹ correspond to the substituents of compound VII.)

55

Table 4 (Compounds in this Table are compounds of the formula VII wherein R⁶ is hydrogen.)

Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ^{2'}	m. p. (°C)
VII-2	H	H	4-F	H	CH ₃	C ₂ H ₅	121- 122
VII-3	H	H	H	H	CH ₃	C ₂ H ₅	102- 102.5
VII-4	H	H	H	H	i-Pr	C ₂ H ₅	85- 85.5
VII-5	6-Cl	H	H	H	CH ₃	C ₂ H ₅	100.5- 101.5
VII-6	6-Cl	H	H	H	i-Pr	C ₂ H ₅	105.5- 106.5
VII-7	H	H	2-F	H	i-Pr	C ₂ H ₅	101.0- 102.0
VII-8	7-Me	H	H	H	i-Pr	C ₂ H ₅	oil
VII-9	H	H	4-Cl	H	i-Pr	C ₂ H ₅	134.0- 136.5
VII-10	H	H	4-OMe	H	i-Pr	C ₂ H ₅	88.0- 89.0
VII-11	H	H	4-Me	H	i-Pr	C ₂ H ₅	108.5- 109.5
VII-12	6-Cl	H	2-Cl	H	i-Pr	C ₂ H ₅	101.0- -103.0
VII-13	H	H	4-CF ₃	H	i-Pr	C ₂ H ₅	117.5- 119.0
VII-14	H	H	3-Me	4-F	i-Pr	C ₂ H ₅	oil
VII-15	H	H	3-Me	5-Me	i-Pr	C ₂ H ₅	oil
VII-16	6-OMe	7-OMe	4-F	H	i-Pr	C ₂ H ₅	96.0- 98.0
VII-17	H	H	4-F	H	C ₂ H ₅	CH ₃	139.0- 139.5
VII-18	H	H	4-F	H	n-Pr	C ₂ H ₅	oil
VII-19	6-Cl	H	4-F	H	i-Pr	C ₂ H ₅	94.5- 95.5
VII-20	H	H	4-F	H	c-Pr	CH ₃	113.5- 116.5
VII-21	H	H	4-OPh	H	i-Pr	C ₂ H ₅	oil
VII-22	6-Cl	8-Cl	4-F	H	i-Pr	C ₂ H ₅	96.0- 98.0
VII-23	6-Cl	H	H	H	Ph	C ₂ H ₅	118.8- -119.5

5	VII-24	6-Cl	H	H	H	c-Pr CH ₃	97.0- 98.5
	VII-25	H	H	4-F	H	sec-Bu CH ₃	oil
	VII-26	6-Me	H	4-F	H	i-Pr C ₂ H ₅	109.0 -111.0
10	VII-27	6-OMe	7-OMe	4-F	H	c-Pr CH ₃	153.0 -153.5

15

VII-8

H-NMR(in CDCl₃) δ ppm :

20 0.92 (t,3H,J=7Hz), 1.41 (d,6H,J=6Hz) 2.47 (s, 3H), 3.27(Heptaplet,1H,J=6Hz) 3.96 (q,2H,J=7Hz), 7.0 -7.8-
(m, 8H)

25

VII-14

H-NMR(in CDCl₃) δ ppm :

1.01 (t,3H,J=7Hz), 1.42 (d,6H,J=6Hz) 2.38 (s, 3H,J=3Hz), 3.25(Heptaplet,1H,J=6Hz) 4.04 (q,2H,J=7Hz),
6.9-8.1(m, 7Hz)

30

VII -15

H-NMR(in CDCl₃) δ ppm :

35 0.97 (t,3H,J=7Hz), 1.43 (d,6H,J=6Hz) 2.29 (s, 3H), 3.25(Heptaplet,1H,J=6Hz) 4.00 (q,2H,J=7Hz), 6.8-8.0-
(m, 7H)

40

VII-18

H-NMR(in CDCl₃) δ ppm :

0.98 (t,3H,J=7Hz), 1.02 (t,3H,J=7Hz) 1.6-2.3(m, 2H), 2.8-3.1(m,2H) 4.03 (q,2H,J=7Hz), 6.9-8.1(m, 8H)

45

VII-21

H-NMR(in CDCl₃) δ ppm :

50 1.03 (t,3H,J=7Hz), 1.41 (d,6H,J=6Hz) 3.25(Heptaplet,1H,J=6Hz) , 4.05(q,2H,J=7Hz), 6.8-8.1(m, 13H)

55

VII-25

H-NMR(in CDCl₃) δ ppm :

0.97 (d,6H,J=6Hz), 2.0-2.6(m,1H) 2.85 (d,2H,J=7Hz), 3.51(s,3H), 6.8-8.1(m, 8H)

5

In the same manner as in Example 1-b, compounds VI-2 to IV-27 were prepared. (In Table 5, R¹, R², R³, R⁴ and R⁵ correspond to the substituents of compound VI.)

10 Tale 5 (Compounds in this Table are compounds of the
formula VI wherein R⁶ is hydrogen.)

15	Compound	R ¹	R ²	R ³	R ⁴	R ⁵	m. p. (°C)
	VI - 2	H	H	p - F	H	CH ₃	—
	VI - 3	H	H	H	H	CH ₃	149-151
20	VI - 4	H	H	H	H	i - Pr	130- 130.5
	VI - 5	6 - C ℓ	H	H	H	CH ₃	139-141
25	VI - 6	6 - C ℓ	H	H	H	i - Pr	168-169
	VI - 7	H	H	2 - F	H	i - Pr	140.5- 142.0
30	VI - 8	7 - Me	H	H	H	i - Pr	155.0- 157.0
	VI - 9	H	H	4 - C ℓ	H	i - Pr	192.0- 195.0
35	VI - 10	H	H	4 - OMe	H	i - Pr	186.0- 188.5
	VI - 11	H	H	4 - Me	H	i - Pr	161.0- 164.0
	VI - 12	6 - C ℓ	H	2 - C ℓ	H	i - Pr	122.0- 124.0
40	VI - 13	H	H	4 - CF ₃	H	i - Pr	183.0- 186.0
	VI - 14	H	H	3 - Me	4 - F	i - Pr	161.0- 162.5
45	VI - 15	H	H	3 - Me	5 - Me	i - Pr	137.0- 138.0
	VI - 16	6 - Me	7 - OMe	4 - F	H	i - Pr	164.0- 165.0
50	VI - 17	H	H	4 - F	H	C ₂ H ₅	141.5- 143.5
	VI - 18	H	H	4 - F	H	n - Pr	146.5- 148.5
55	VI - 19	6 - C ℓ	H	4 - F	H	i - Pr	171.0- 172.0

VI-20	H	H	4-F	H	c-Pr	120-126
VI-21	H	H	4-OPh	H	i-Pr	153.0- 154.0
VI-22	6-Cl	8-Cl	4-F	H	i-Pr	98.5-103
VI-23	6-Cl	H	H	H	Ph	171.5- 172.5
VI-24	6-Cl	H	H	H	c-Pr	84.0- 86.0
VI-25	H	H	4-F	H	sec-Bu	119.0- 121.0
VI-26	6-Me	H	4-F	H	i-Pr	160.0- 161.5
VI-27	6-OMe	7-OMe	4-F	H	c-Pr	162.0- 163.0

In the same manner as in Example 1-c, compounds V-2 to V-27 were prepared. (In Table 6, R¹, R², R³, R⁴ and R⁵ correspond to the substituents of compound of V.)

Table 6 (Compounds in this Table are compounds of the formula V wherein R⁶ is hydrogen.)

Compound	R ¹	R ²	R ³	R ⁴	R ⁵	m. p. (°C)
V-2	H	H	p-F	H	CH ₃	125-128
V-3	H	H	H	H	CH ₃	143-146
V-4	H	H	H	H	i-Pr	92-93
V-5	6-Cl	H	H	H	CH ₃	220-222

	V-6	6-Cl	H	H	H	i-Pr	140-140.5
	V-7	H	H	2-F	H	i-Pr	121.5- 124.0
5	V-8	7-Me	H	H	H	i-Pr	105.1- 109.2
	V-9	H	H	4-Cl	H	i-Pr	147.0- 147.8
10	V-10	H	H	4-OMe	H	i-Pr	135.6- 136.8
	V-11	H	H	4-Me	H	i-Pr	119.4- 120.4
15	V-12	6-Cl	H	2-Cl	H	i-Pr	105.8- 106.9
	V-13	H	H	4-CF ₃	H	i-Pr	163.7- 164.2
	V-14	H	H	3-Me	4-F	i-Pr	161.1- 108.1
20	V-15	H	H	3-Me	5-Me	i-Pr	120.8- 122.3
	V-16	6-OMe	7-OMe	4-F	H	i-Pr	164.4- 165.2
25	V-17	H	H	4-F	H	C ₂ H ₅	143.1- 144.2
	V-18	H	H	4-F	H	n-Pr	150.2- 155.3
	V-19	6-Cl	H	4-F	H	i-Pr	164.5- 165.3
30	V-20	H	H	4-F	H	c-Pr	150.1- 151.6
	V-21	H	H	4-OPh	H	i-Pr	106.9- 107.7
35	V-22	6-Cl	8-Cl	4-F	H	i-Pr	135.0- 135.7
	V-23	6-Cl	H	H	H	Ph	174.8- 175.3
40	V-24	6-Cl	H	H	H	c-Pr	157.5- 158.0
	V-25	H	H	4-F	H	sec-Bu	125.0- 126.5
	V-26	6-Me	H	4-F	H	i-Pr	155.0- 157.0
45	V-27	6-OMe	7-OMe	4-F	H	c-Pr	200.0- 200.5

50 In the same manner as in Example 1-d, compounds IV-2 to IV-6 were prepared. (In Table 7, R¹, R², R³, R⁴ and R⁵ correspond to the substituents of compound of IV.)

55

Table 7

(Compounds in this Table are compounds of the formula IV wherein R⁶ is hydrogen.)

Compound	R ¹	R ²	R ³	R ⁴	R ⁵	m. p. (°C)
IV - 2	H	H	4-F	H	CH ₃	177-179
IV - 3	H	H	H	H	CH ₃	-
IV - 4	H	H	H	H	i-Pr	-
IV - 5	6-Cl	H	H	H	CH ₃	-
IV - 6	6-Cl	H	H	H	i-Pr	-

In the same manner as in Example 1-a, compounds III-2 to III-27 were prepared. (In Table 8, R¹, R², R³, R⁴ and R⁵ correspond to the substituents of compound III.)

Table 8 (Compounds in this Table are compounds of the formula III wherein R⁶ is hydrogen.)

Compound	R ¹	R ²	R ³	R ⁴	R ⁵	m. p. (°C)
III - 2	H	H	4-F	H	CH ₃	194-196
III - 3	H	H	H	H	CH ₃	170-171.5
III - 4	H	H	H	H	i-Pr	107-108.5
III - 5	6-Cl	H	H	H	CH ₃	192-194
III - 6	6-Cl	H	H	H	i-Pr	125.5-127
III - 7	H	H	2-F	H	i-Pr	80.1-80.2
III - 8	7-Me	H	H	H	i-Pr	121.1-122.3
III - 9	H	H	4-Cl	H	i-Pr	148.0-149.1
III - 10	H	H	4-OMe	H	i-Pr	137.4-140.1
III - 11	H	H	4-Me	H	i-Pr	111.6-113.1
III - 12	6-Cl	H	2-Cl	H	i-Pr	83.8-84.5
III - 13	H	H	4-CF ₃	H	i-Pr	126.2-128.8

	III -14	H	H	3-Me	4-F	i-Pr	124.8- 126.4
5	III -15	H	H	3-Me	5-Me	i-Pr	117.6- 120.3
	III -16	6-OMe	7-OMe	4-F	H	i-Pr	147.8- 150.9
	III -17	H	H	4-F	H	C ₂ H ₅	124.3- 128.5
10	III -18	H	H	4-F	H	n-Pr	117.8- 121.5
	III -19	6-Cℓ	H	4-F	H	i-Pr	135.2- 135.9
15	III -20	H	H	4-F	H	c-Pr	141.3- 144.1
	III -21	H	H	4-OPh	H	i-Pr	oil
	III -22	6-Cℓ	8-Cℓ	4-F	H	i-Pr	117- 122
20	III -23	6-Cℓ	H	H	H	Ph	142.8- 144.3
	III -24	6-Cℓ	H	H	H	c-Pr	161.0- 161.5
25	III -25	H	H	4-F	H	sec-Bu	78.0- 81.0
	III -26	6-Me	H	4-F	H	i-Pr	137.0- 137.5
30	III -27	6-OMe	7-OMe	4-F	H	c-Pr	189.5- 191.0

35
III-22

H-NMR(in CDCl₃) δ ppm :
 1.40(d,6H,J=7Hz), 3.44(Heptaplet,1H,J=7Hz) 5.93(dd,1H,J=8Hz,J=16Hz), 6.8-8.1(m,14H) 9.34-
 40 (d,1H,J=8Hz)

In the same manner as in Example 1-f, compounds II-2 to II-27 were prepared. (In Table 9, R¹, R², R³, R⁴ and R⁵ correspond to the substituents of compound II.)

45

50

55

Table 9 (Compounds in this Table are compounds of the formula of II wherein R⁶ is hydrogen.)

Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ^{1,2}	m. p. (°C)
II-2	H	H	p-F	H	CH ₃	C ₂ H ₅	oil
II-3	H	H	H	H	CH ₃	C ₂ H ₅	105 -106
II-4	H	H	H	H	i-Pr	C ₂ H ₅	88.5 -90.5
II-5	6-Cl	H	H	H	CH ₃	C ₂ H ₅	77-82
II-6	6-Cl	H	H	H	i-Pr	C ₂ H ₅	96-98
II-7	H	H	2-F	H	i-Pr	C ₂ H ₅	oil
II-8	7-Me	H	H	H	i-Pr	C ₂ H ₅	68.5- 74.0
II-9	H	H	4-Cl	H	i-Pr	C ₂ H ₅	91.0 -94.0
II-10	H	H	4-OMe	H	i-Pr	C ₂ H ₅	78.0 -78.5
II-11	H	H	4-OMe	H	i-Pr	C ₂ H ₅	75.0 -78.0
II-12	6-Cl	H	2-Cl	H	i-Pr	C ₂ H ₅	oil
II-13	H	H	4-CF ₃	H	i-Pr	C ₂ H ₅	78.0 -83.0
II-14	H	H	3-Me	4-F	i-Pr	C ₂ H ₅	66.0 -71.0
II-15	H	H	3-Me	5-Me	i-Pr	C ₂ H ₅	oil

	II -16	6-OMe	7-OMe	4-F	H	i-Pr	C ₂ H ₅	83.0
								-90.0
	II -17	H	H	4-F	H	C ₂ H ₅	C ₂ H ₅	94.0
5								-97.0
	II -18	H	H	4-F	H	n-Pr	C ₂ H ₅	oil
	II -19	6-Cl	H	4-F	H	i-Pr	C ₂ H ₅	111.0-
								113.5
10	II -20	H	H	4-F	H	c-Pr	C ₂ H ₅	91.0
								-93.0
	II -21	H	H	4-OPh	H	i-Pr	C ₂ H ₅	121.0-
								125.0
15	II -22	6-Cl	8-Cl	4-F	H	i-Pr	C ₂ H ₅	oil
	II -23	6-Cl	H	H	H	Ph	C ₂ H ₅	oil
	II -24	6-Cl	H	H	H	c-Pr	C ₂ H ₅	69.0
20	II -25	H	H	4-F	H	sec-Bu	C ₂ H ₅	oil
								-71.0
	II -26	6-Me	H	4-F	H	i-Pr	C ₂ H ₅	oil
25	II -27	6-OMe	7-OMe	4-F	H	c-Pr	C ₂ H ₅	oil

30 II-7

H-NMR(in CDCl₃) δ ppm :

1.21(t,3H,J=7Hz), 1.32(d,6H,J=6Hz) 2.2-2.4(m,2H), 2.5-2.7(m,1H) 3.28 (s, 1H), 3.34(Heptaplet,1H,J=6Hz)
 4.08(q,2H,J=7Hz), 4.3-4.6(m, 1H) 5.28(dd,1H,J=1.5Hz,J=15Hz), 6.53(dd, 1H,J=1.5Hz,J=15Hz), 6.9-8.0-
 35 (m,8H)

40 II-12

H-NMR(in CDCl₃) δ ppm :

1.25(t,3H,J=7Hz), 1.33(d,6H,J=6Hz) 2.2-2.4(m,2H), 2.5-2.8(m,1H) 3.32(s, 2H), 3.38(Heptaplet,1H,J=6Hz)
 4.13(q,2H,J=7Hz), 4.2-4.6(m, 1H) 5.34(dd,1H,J=6Hz, J=15Hz), 6.53(dd,1H,J=1.5Hz,J=15Hz), 7.0-8.0-
 45 (m,7H)

II-15

50 H-NMR(in CDCl₃) δ ppm :

1.23(t,3H,J=7Hz), 1.35(d,6H,J=6Hz) 2.2-2.4(m,2H), 2.31(s,6H) 2.6-2.8(m,1H), 3.32(s, 2H) 3.35-
 (Heptaplet,1H,J=6Hz),4.12(q,2H,J=7Hz) 4.3-4.7(m,1H), 5.30(dd,1H,J=6Hz,J=16Hz) 6.51-
 (dd,1H,J=1Hz,J=16Hz), 6.7-8.0(m,7H)

55

II-18

H-NMR(in CDCl₃) δ ppm :

1.00(t,3H,J = 7Hz), 1.26(t,3H,J = 7Hz) 1.6-2.3(m,2H), 2.42(d,2H,J = 6Hz) 2.6-3.2(m,3H), 3.35(s,2H) 4.11-
 5 (q,2H,J = 7Hz), 4.3-4.7(m, 1H) 5.27(dd,1H,J = 6Hz,J = 16Hz) 6.46(dd,1H,J = 1.5Hz,J = 16Hz), 6.9-8.0(m,8H)

II-22

10

H-NMR(in CDCl₃) δ ppm :

1.26(t,3H,J = 7Hz), 1.33(d,6H,J = 6Hz) 2.43(d,2H,J = 6Hz), 2.6-2.9(m,1H) 3.36(s, 2H), 3.44-
 (Heptaplet,1H,J = 6Hz) 4.13(q,2H,J = 7Hz), 4.3-4.7(m, 1H) 5.30(dd,1H,J = 6Hz,J = 16Hz), 6.53-
 15 (dd,1H,J = 1.5Hz,J = 16Hz), 7.0-7.6(m,6H)

II-23

20 H-NMR(in CDCl₃) δ ppm :

1.23(t,3H,J = 7Hz), 2.21(d,2H,J = 6Hz) 2.4-2.6(m,1H), 3.25(s,2H) 4.09(q,2H,J = 7Hz), 4.1-4.4(m, 1H) 5.08-
 (dd,1H,J = 6Hz,J = 16Hz), 6.26(dd,1H,J = 1.5Hz,J = 16Hz), 7.0-8.0 (m,13H)

25

II-25

H-NMR(in CDCl₃) δ ppm :

0.96(d,6H,J = 6Hz), 1.26(t,3H,J = 7Hz), 1.8-2.4(m,1H), 2.43(d,2H,J = 6Hz), 2.6-2.9(m,1H), 2.88(d,2H,J = 7Hz),
 30 3.36(s,2H), 4.14(q,2H,J = 7Hz), 4.3-4.7(m, 1H), 5.0-5.5(m,1H), 6.3-6.7(m,1H), 6.9-8.1(m,8H)

II-26

35

H-NMR(in CDCl₃) δ ppm :

1.25(t,3H,J = 7Hz), 1.32(d,6H,J = 6Hz), 2.32(s,3H), 2.39(d,2H,J = 7Hz), 2.6-3.1(m,1H), 3.36(s, 2H), 3.41-
 (Heptaplet,1H,J = 6Hz), 4.11(q,2H,J = 7Hz), 4.3-4.7(m, 1H), 5.0-5.5(m,1H), 6.3-6.7(m,1H), 6.8-7.9(m,7H)

40

II-27

H-NMR(in CDCl₃) δ ppm :


0.8-1.5(m,4H), 1.26(t,3H,J = 7Hz), 2.0-2.9(m,4H), 3.42(s,2H), 3.71(s,3H), 4.00(s,3H), 4.20(q,2H,J = 7Hz), 4.4-
 45 4.8(m,1H), 5.3-5.8(m,1H), 6.4-6.9(m,1H), 6.58(s,1H), 7.0-7.5(m,5H)

In the same manner as in Example 1-g, compounds I-12 to I-127 were prepared.

50

55

Table 10



Compound	R ¹	R ²	R ³	R ⁴	R ⁵	R ¹²	m.p. (°C) Mass spectrum
I - 12	H	H	4-F	H	CH ₃	C ₂ H ₅	oil M/e 423, 292 264, 249
I - 13	H	H	H	H	CH ₃	C ₂ H ₅	92-105
I - 14	H	H	H	H	i-Pr	C ₂ H ₅	97-100
I - 15	6-Cl	H	H	H	CH ₃	C ₂ H ₅	oil

EP 0 304 063 A2

	I -16	6-Cl	H	H	H	i-Pr C ₂ H ₅	oil
	I -17	H	H	2-F	H	i-Pr C ₂ H ₅	oil
5	I -18	7-Me	H	H	H	i-Pr C ₂ H ₅	oil
	I -19	H	H	4-Cl	H	i-Pr C ₂ H ₅	98-104
10	I -110	H	H	4-OMe	H	i-Pr C ₂ H ₅	94-98
	I -111	H	H	4-Me	H	i-Pr C ₂ H ₅	79-85
15	I -112	6-Cl	H	2-Cl	H	i-Pr C ₂ H ₅	oil
	I -113	H	H	4-CF ₃	H	i-Pr C ₂ H ₅	117-128
	I -114	H	H	3-Me	4-F	i-Pr C ₂ H ₅	85-92
20	I -115	H	H	3-Me	5-Me	i-Pr C ₂ H ₅	oil
	I -116	6-OMe	7-OMe	4-F	H	i-Pr C ₂ H ₅	gum
25	I -117	H	H	4-F	H	C ₂ H ₅ C ₂ H ₅	oil
	I -118	H	H	4-F	H	n-Pr C ₂ H ₅	oil
	I -119	6-Cl	H	4-F	H	i-Pr C ₂ H ₅	79-82
30	I -120	H	H	4-F	H	c-Pr C ₂ H ₅	100-104
	I -121	H	H	4-OPh	H	i-Pr C ₂ H ₅	oil
35	I -122	6-Cl	8-Cl	4-F	H	i-Pr C ₂ H ₅	133-143
	I -123	6-Cl	H	H	H	Ph C ₂ H ₅	gum
40	I -124	6-Cl	H	H	H	c-Pr C ₂ H ₅	oil
	I -125	H	H	4-F	H	sec-Bu C ₂ H ₅	oil
45	I -126	6-Me	H	4-F	H	i-Pr C ₂ H ₅	oil
	I -127	6-OMe	7-OMe	4-F	H	c-Pr C ₂ H ₅	gum

50

55

EP 0 304 063 A2

I-17

H-NMR(in CDCl_3) δ ppm :

1.29(t,3H,J=7Hz), 1.40(d,6H,J=6Hz) 1.4-1.7(m,2H), 2.3-2.5(m,2H) 2.9-3.2(m,1H), 3.49(Heptaplet,1H,J=6Hz)
5 3.5-3.8(m,1H), 3.9-4.5(m,2H) 4.20(q,2H,J=7Hz), 5.2-5.7(m, 1H) 6.5-6.9(m,1H), 7.0-8.2(m,8H)

I-18

70

H-NMR(in CDCl_3) δ ppm :

1.0-1.4(m,2H), 1.31(t,3H,J=7Hz) 1.39(d,6H,J=6Hz), 2.3-2.5(m,2H) 2.52(s,3H), 3.1-3.4(m,1H) 3.48-
(Heptaplet,1H,J=6Hz),3.5-3.8(m,1H) 3.8-4.1(m, 1H), 4.20(q,2H,J=7Hz) 4.2-4.5(m,1H), 5.2-5.6(m,1H) 6.4-6.8-
(m,1H), 7.0-8.0(m,8H)

75

I-19

20 H-NMR(in CDCl_3) δ ppm :

1.29(t,3H,J=7Hz), 1.38(d,6H,J=6Hz) 1.4-1.8(m,2H), 2.3-2.5(m,2H) 3.2-3.4(m,1H), 3.49(Heptaplet,1H,J=6Hz)
3.6-3.8(m,1H), 3.9-4.2(m,1H) 4.20(q,2H,J=7Hz), 4.3-4.5(m,1H) 5.2-5.5(m,1H), 6.5-6.8(m,1H) 7.0-8.2(m,8H)

25

I-110

H-NMR(in CDCl_3) δ ppm :

1.29(t,3H,J=7Hz), 1.40 (d,6H,J=6Hz) 1.5-1.6(m,2H), 2.3-2.5(m,2H) 2.8-3.0(m,1H), 3.4-3.6(m,1H) 3.52-
30 (Heptaplet,1H,J=6Hz),3.88(s,3H) 3.9-4.1(m,1H), 4.20(q,2H,J=7Hz) 4.3-4.5(m,1H), 5.3-5.5(m,1H) 6.5-6.7-
(m,1H), 6.9-8.1(m,8H)

35 I-111

H-NMR(in CDCl_3) δ ppm :

1.30(t,3H,J=7Hz), 1.3-1.5(m,2H) 1.39(d,6H,J=6Hz), 2.3-2.5(m,2H) 2.43(s,3H), 2.8-3.0(m,1H) 3.50-
(Heptaplet,1H,J=6Hz),3.5-3.7(m,1H) 3.9-4.2(m,1H), 4.19(q,2H,J=7Hz) 4.2-4.5(m,1H), 5.2-5.6(m,1H) 6.4-6.8-
40 (m,1H), 6.9-8.2(m,8H)

I-112

45

H-NMR(in CDCl_3) δ ppm :

1.30(t,3H,J=7Hz), 1.3-1.6(m,2H) 1.37(d,6H,J=6Hz), 2.3-2.5(m,2H) 2.9-3.2(m,1H), 3.47(Heptaplet,1H,J=6Hz)
3.5-3.8(m,1H), 3.9-4.1(m,1H) 4.19(q,2H,J=7Hz), 4.2-4.5(m, 1H) 5.3-5.7(m,1H), 6.5-6.8(m,1H) 7.1-8.1(m,7H)

50

I-113

H-NMR(in CDCl_3) δ ppm :

1.0-1.3(m,2H), 1.30(t,3H,J=7Hz) 1.40(d,6H,J=6Hz), 2.3-2.4(m,2H) 3.3-3.5(m,1H), 3.49(Heptaplet,1H,J=6Hz)
55 3.6-3.7(m,1H), 3.9-4.1(m,1H) 4.18(q,2H,J=7Hz), 4.2-4.5(m,1H) 5.1-5.5(m, 1H), 6.5-6.8(m,1H) 7.2-8.2(m,8H)

PATENT ABSTRACTS OF JAPAN

(11)Publication number : 08-127585

(43)Date of publication of application : 21.05.1996

(51)Int.Cl.

C07F 7/18
// C07D405/06
C07M 7:00

(21)Application number : 06-276395

(71)Applicant : UBE IND LTD
NISSAN CHEM IND LTD

(22)Date of filing : 10.11.1994

(72)Inventor : HARADA KATSUMASA
MATSUSHITA AKIO
KAWACHI YASUHIRO
SASAKI HIROSHI

(30)Priority


Priority number : 06212960 Priority date : 06.09.1994 Priority country : JP

(54) OPTICALLY ACTIVE 3-OXY-5-OXO-5-HEPTENOIC ACID DERIVATIVE AND ITS PRODUCTION

(57)Abstract:

PURPOSE: To obtain a new compound useful as a synthetic intermediate for a blood cholesterol lowering agent.

CONSTITUTION: A methyl (3R,6E)-3-[(tert-butyl dimethylsilyl)oxy]-7-[2'-cyclopropyl-4'-(4"-fluorophenyl) quinolin-3'-yl]-5-oxo-6-heptenoate of the formula. The compound of the formula is obtained in good yield for a short time by reacting (A) 2-cyclopropyl-4-(4'- fluorophenyl)-3-quinolinecarboxyaldehyde with (B) methyl (R)-3-tert-butyl dimethylsilyloxy-6-dimethoxyphosphinyl-5-oxo-hexanoate in the presence of potassium carbonate in an aliphatic alcohol solvent. Furthermore, the aliphatic alcohol solvent preferably mainly contains a secondary or a tertiary aliphatic alcohol and further contains about 30-20000ppm moisture.



LEGAL STATUS

[Date of request for examination] 29.03.2001

[Date of sending the examiner's decision of rejection]

[Kind of final disposal of application other than the examiner's decision of rejection or application converted registration]

[Date of final disposal for application]

[Patent number]

[Date of registration]

[Number of appeal against examiner's decision
of rejection]

[Date of requesting appeal against examiner's
decision of rejection]

[Date of extinction of right]

Copyright (C); 1998,2003 Japan Patent Office